EU 584

18 APR 1990

ISSN 0013-8908

THE ENTOMOLOGIST'S MONTHLY MAGAZINE

World List Abbreviation, 4th ed. (1963): *Entomologist's mon. Mag.*

EDITED BY
Prof. Sir Richard SOUTHWOOD, D.Sc., F.R.S., F.R.E.S.
M. G. MORRIS, M.A., Ph.D., F.I.Biol., F.L.S., F.R.E.S.
K. G. V. SMITH, C.Biol., F.I.Biol., F.L.S., F.Z.S., F.R.E.S.
J. W. ISMAY, B.Sc., Ph.D., F.R.E.S.
P. M. HAMMOND, M.A., F.R.E.S.
W. R. DOLLING, B.Sc., F.R.E.S.

VOLUME 125

'J'engage donc tous à éviter dans leurs écrits toute personnalité, toute allusion dépassant les limites de la discussion la plus sincère et la plus courtoise.' — *Laboulbène.*

THE ENTOMOLOGIST'S MONTHLY MAGAZINE
BRIGHTWOOD
BRIGHTWELL, WALLINGFORD, OXON OX10 0QD
1989

Published by Gem Publishing Company

CONTENTS

CONTRIBUTORS iii
GENERAL INDEX iv
SPECIAL INDEX –
ARACHNIDA vii
COLEOPTERA vii
COLLEMBOLA ix
DERMAPTERA ix
DIPTERA ix
HEMIPTERA xi
HYMENOPTERA xi
ISOPTERA xii
LEPIDOPTERA xii
MANTODEA xii
MECOPTERA xii
NEUROPTERA xii
ODONATA xii
ORTHOPTERA xiii
PHTHIRAPTERA xiii
PSOCOPTERA xiii
SIPHONAPTERA xiii
THYSANURA xiii
GROUPS OTHER THAN INSECTA AND ARACHNIDA xiii
ERRATA AND CORRIGENDA xiii

DATES OF PUBLICATION

January/February/March/April	31st March 1989
May/June/July/August	31st July 1989
September/October/November/December	30th November 1989

INDEX

This index covers the year 1989 (Vol. **125**). Items occurring on the wrappers are indicated by Roman numerals. Prepared by Mrs A.V. Smith.

INDEX TO CONTRIBUTORS

Ackland, D.M. 211
Agarwal, M.L., 31
Alexander, K.N.A., 48, 84, 106, 127
Allen, A.A., 3, 12, 24, 51, 79, 121, 132, 137, 138, 140, 181, 182
Allen, G.W., 95, 103
Anderson, R., 4
Archer, M.E., 103, 143, 206, 232
Blackman, R.L., 236
Bouček, Z., 85
Brindle, A., 13
Brownbridge, M., 123
Bunn, D.S., 117, 133
Carter, D.J., 80, 94, 128
Chainey, J.E., 141
Clements, D.K., 153
Coldwell, J.D., 250
Collier, M., 167
Collins, G.B., 250
Cooter, J., 242, 244
Cox, M.L., 87, 175
Crowson, R.A., 129, 210
Dike, M.C., 73
Disney, R.H.L., 45, 81, 169, 183
Dowie, M.T., 237
Edwards, R., 106
Else, G.R., 95, 106, 253
Ford, T.H., 139
Foster, G.N., 71
Fowler, J.A., 118
Fowles, A.R., 177
Fox, A.D., 151
Freeman, P., 37
Godfray, A., 252
Goldie-Smith, E.K., 107
Graham, M.W.R. de V., 85, 231
Greenwood, S., 36
Hammond, P.M., 95
Hanson, H.M., 245
Harris, K.M., xi
Hawkins, R.D., 174, 205, 241
Heath, R.V., 18
Henshaw, D., 83
Horsfield, D., 182, 230, 243, 249
Howse, D., 83
Hutson, A.M., 122, 176
Jefferies, D.J., 245
Kapoor, V.C., 31
Lane, S.A., 36, 127
Lansbury, I., 63
Laurence, B.R., 89
Leather, S.R., 17
Liston, A.D., 62, 168
Littlewood, F.M., 256
Littlewood, P.M.H., 233, 256
Luff, M.L., 118
Luna Dias, A.P.A., 44
Lyal, C.H.C., 24
Lyszkowski, R.M., 8
Majerus, M.E.N., 177
Meharg, M., 4
Miles, S., 168
Morris, M.G., 25, 30, 128, 168
Mulcahy, R.L., 237
O'Connor, J.P., 206
Owen, J.A., 156
Pearce, M.J., 207
Pennington, T.H., 132
Pitkin, B., 52
Pont, A.C., 243
Pontin, A.J., 4
Quicke, D.L.J., 9, 19, 119, 199
Read, R.W.J., 30, 117
Reynolds, W.J., 251
Rotheray, G.E., 5, 8, 174
Schroth, M., 169
Scoble, M.J., 239
Selman, B.J., 123
Shaw, M.R., 174
Shelley, A.J., 44
Sinclair, M., 198
Smith, K.G.V., 43, 52, 61, 94, 95, 105, 122, 138, 141, 149, 210, 235, 253
Southwood, Prof. Sir Richard, 1
Vardy, C.R., 105
Verdcourt, B., 82, 132
Walker, A., 19
Wanless, F.R., 88
Weaving, A.J.S., 157
White, I.M., 53
Whitehead, P.F., 49, 79, 118, 150, 187, 198, 236
Williams, T., 72
Wilson, M.R., 135
Wright, A., 127

GENERAL INDEX

Adaptation of aphid eggs to specific ant hosts, 4
Africa, some new species of *Atherigona* Rondani (Dipt., Muscidae) from, 73
Australasian Chalcidoidea (Hymenoptera): Errors and omissions, 85
Bardsey Island (Ynes Enlli), meniscus midges, (Dipt., Dixidae) on, 118
Bark, *Medetera* (Dipt., Dolichopodidae) cocoons, under, 174
Bedfordshire, *Otiorhynchus scaber* (L.) (Col., Curculionidae) in – a correction, 132
Bees Wasps and Ants, recording scheme, xi
Belated records of three rare Ptiliidae (Col.), 138
Bird cherry (*Prunus padus*), *Phytodecta pallida* (L.) (Col., Chrysomelidae), a new insect record for, 17
Bird and mammal nests and mammal runs, burrows and droppings, E.B. Basden's collection of Diptera from, 5
Blackflies (Dipt., Simuliidae), first report of man eating, 44
Breckland and Suffolk Sandings, notable wasps and bees (Hym., Aculeata), in, 232
Britain, a new insect record for, 17; a further record of *Lutridia exilis* (Mitzsch) (Phthiraptera, Trichodectidae) in Britain, with notes on the presence and absence of lice on, 245; a new insect record for, 17; Anthomyiidae (Dipt.) new to, with a description of new species of *Botanophila* Lioy, 211; *Dolichovespula saxonica,* Fabricius 1793 (Hym., Vespidae) in, with a key to British *Dolichovespula,* 103; *Euderomphale chelidonii* (Erdös) (Hym., Eulophidae); first record in, 72; *Medina separata* (Meigen) (Dipt., Tachinidae), new to, 139; occurrence of *Leopoldius brevirostris* (Germar) (Dipt., Conopidae) in, with notes on the distinction from *L. signatus* (Wiedemann), 153; *Orsillus depressus* Dallas (Hem., Lygaeidae), an arboreal groundbug new to, 241; *Spilogona trigonata* (Zetterstedt, 1838) (Dipt., Muscidae) new to, 243; the re-discovery of the 5-spot ladybird (*Coccinella 5-punctata* L.) (Col., Coccinellidae) in, 177; thirty years of *Eriozona syrphoides* (Fallén) (Dipt., Syrphidae) in, 132
British insect fauna, some recent additions to the, 95
Buckinghamshire, *Hemerobius marginatus* Stephens (Neurop., Hemerobiidae) in, 82
Cannibalism in *Mitopus morio* (F.) (Arach., Opiliones) observed in the field, 256
Cape figwort, *Cionus scrophulariae* (L.) (Col., Curculionidae) breeding in, in England, 150
Carcinophora americana (Beauvois) and related species (Derm., Anisolabididae), 13
Cave-dwelling fly, *Speolepta leptogaster* Winn. (Dipt., Mycetophilidae) in Central London, 89
Centaurea solstitialis L., a new species of *Terellia* Robineau-Desvoidy associated with, and a revision of the *Terellia virens* species group (Dipt., Tephritidae), 53
Cheshire, *Leistus rufomarginatus* (Dufts.) (Col., Carabidae) from, 18
Conifer plantations, *Loberus pratorum* (Fallén) (Hym., Tenthredinidae) locally abundant in, 168
corrections — 105, 121, 132
Croydon, *Arhopalus rusticus* (L.) (Col., Cerambycidae) in South, 250
Cultures of leatherjackets (*Tipula* spp.) (Dipt., Tipulidae), improvements in rearing laboratory, 123
Cumbria, *Ceutorhynchus euphorbiae* Brisout (Col., Curculionidae) in west, 117; *Ceutorhynchus mixtus* Mulsant & Rey (Col. Curculionidae) in west, 30
Defecation behaviour, differences in, between *Dolichovespula sylvestris* (L.) and *D. norwegica* (F.) (Hym.), 117
Densities, high, of cranefly larvae (Dipt., Tipulidae), 8
Eggs of *Dixella aestivalis* Meigen, and brief comparisons with eggs of three other species of *Dixella* (Dipt., Dixidae), 107
English records of *Dolichovespula media* (Retzius) (Hym., Vespidae), 106
European Agromyzidae (Diptera), revised status of eight species of, described both as larvae and adults, and four new names for homonyms, 83
Falkland Islands (Malvinas), two new species of *Megaselia* (Dipt., Phoridae) from the, and a replacement name in the genus, 183
Feeding behaviour of *Panorpa communis* (L.) (Mecopt., Panorpidae), 36
First quarter of the second century of the *E.M.M.*, the, 1
Foodplant, *Miarus micros* (Germ.) (Col., Curculionidae): a problem of, 3
France, a new species of *Phora* (Dipt., Phoridae) from, 81
Further record of *Placochilus seladonicus* (Fallén) (Hem., Miridae), 205
Gilchrist Expedition Award, xv

Gloucestershire, *Anisodactylus binotatus* (F.) (Col., Carabidae) in, 79; beetles from unvegetated wet sediments at Bishop's Cleeve, 150; *Meloe rugosus* Marsham (Col., Meloidae) in, 127
Gwent, *Oncomera femorata* (F.) (Col., Oedomeridae) hibernating near Chepstow, with a note of associated invertebrates, 236
Hampshire, *Andrena hattorfiana* (F.) (Hym., Andrenidae) in, 168; *Colon dentipes* (Sahlberg) (Col., Leiodidae) in south, 242
Hering Memorial Research Fund, v
Hertfordshire, *Melanophthalma suturalis* (Mann.) (Col., Lathridiidae) in Bucks and, 140; *Tomoxia bucephala* Costa (Mordellidae), *Bitoma crenata* (F.) (Colydidae) and other local Coleoptera from Whipendell Wood, 252
Host records for genera and species of Braconinae (Hym., Braconidae), further new, 199
Identity of four unique British specimens of *Gymnetron* (Col., Curculionidae), 25
India, new Tephritidae (Diptera) from, 31
Indo-Australian genus of Braconinae (Hym., Braconidae) parasitic on Hispidae (Col.), a new, 19
Infestation of *Archaeopsylla erinacei erinacei* (Bouché) (Siphonaptera, Pulicidae) with hypopi of *Acarus nidicolous* (Acari: Astigmata), 237
Inter- and intraspecific usurpation in *Dolichovespula* (Hym., Vespidae), 133
Inverness-shire, *Cephalcia lariciphila* (Wachtl) (Hym., Pamphiliidae) in, and the significance of conifer sawflies on mountain summits, 62
Ireland, a third site for *Trechus rivularis* (Gyll.) (Col., Carabidae) in, 4; notes on Irish *Temnostethus* (Hem.-Het., Anthocoridae) in the National Museum of, 206
Isle of Man, 1987, further new Coleoptera from the, 118
Kent, *Dilta hibernica* (Carpenter) (Thysanura, Machilidae) in, 48; *Lamprohiza splendidula* (L.) (Col., Lampyridae) taken in 1884, 182; *Stenus formicetorum* Mann. (Col., Staphylinidae), plentiful in a suburban straw-stack, apparently new to, 137
London, the ecology of a cave-dwelling fly, *Speolepta leptogaster* Winn. (Dipt., Mycetophilidae) in Central, 89
Mammal nests, runs, burrows and droppings and bird nests, E.B. Basden's collection of Diptera from, 5
Man eating blackflies (Dipt., Simuliidae), first report of, 44
Monmouthshire, *Philonthus rotundicollis* (Ménétriés) (Col., Staphylinidae) in, 244
Neotropical genus and species of Coeloidini (Hym., Braconidae, Braconinae) a new, 9
Nesting behaviour and prey of two Southern African species of *Stizus* (Hym., Sphecidae, Nyssoninae), notes on, 157
New World species of Braconinae (Hym., Braconidae), reclassification of some, 119
Niche differences in the nesting sites of British social wasps (Hym., Vespinae), 143
Nigeria, two new species of *Megaselia* (Dipt., Phoridae) reared from termite-mound fungus in, 45
Non-British species of *Anapausis* (Dipt., Scatopsidae), some, 37
Norfolk, *Rhynchaenus testaceus* (Müller) (Col., Curculionidae) in, 167
OBITUARY – Sir Christopher Howard Andrewes, 253
Ommatidial size-differentiation, observations on *Megaselia persecutrix* Schmitz (Dipt., Phoridae) and the significance of, 169
Outbreak of *Phyllodecta laticollis* Suffr. (Col., Chrysomelidae): with a comment on the present distribution and incidence of *P. vulgatissima* (L.), an, 24
Oviposition behaviour in *Somatochlora metallica* van der Linden (Odonata), 151
Oxford record of *Sepedophilus testaceus* (F.) (Col., Staphylinidae), an, 12
Perthshire, a record of *Ernoneura argus* (Zett.) (Dipt., Scathophagidae) from, 230
Recent records of *Cassida nebulosa* L. (Col., Chrysomelidae), 168
REVIEWS – **Agrios, G.N.,** 1988, Plant Pathology, xi; **Bachli, G. & Buria, H.,** Diptera Drosophilidae (*Insecta Helvetica* **7**), 52; **Barbosa, P. & Schultz, J.C.** (eds), 1987, Insect Outbreaks, 236; **Bouček, Z.,** 1988, Australasian Chalcidoidea (Hymenoptera) . . ., 85; **Bovey, P.,** 1987. Coleoptera, Scolytidae, Platypodidae (*Insecta Helvetica* **6**) 87; **Costa, A., Vanin, S.A. & Casari-Chen, S.A.,** 1988, Larvas de Coleoptera do Brasil, 210; **Cranston, P.S., Ramsdale, C.D., Snow, K.R. & White, G.B.,** 1987, Keys to the adults, male hypopygia, fourth instar larvae and pupae of the British mosquitoes (Culicidae) . . ., 52; **Eaton, J.L.,** 1988, Lepidopteran Anatomy, 239; **Ferrar, P.,** 1987, A Guide to the Breeding Habits and Immature Stages of Diptera Cyclorrhapha, 61; **Foelix, R.F.,** 1987, Biology of Spiders, 88; **Hansen, M.,** 1987, The Hydrophiloidea (Coleoptera) of Fennoscandia and Denmark (*Faun. Ent. Scand.* **18**), 175; **Harrison, F. & Sterling, J.,** 1988, Butterflies and Moths of Derbyshire, 128; **Holt, V.M.,** 1988, Why Not Eat

Insects?, 43; **Jobling, B.,** 1987, Anatomical Drawings of Biting Flies, 122; **Kudrna, P.,** 1986, Butterflies of Europe, Vol. 8, Aspects of the Conservation of Butterflies in Europe, 80; **Marshall, J.A. & Haes, E.C.M.,** 1987, Grasshoppers and Allied Insects of Great Britain and Ireland, 251; **Pape, T.,** 1987, The Sarcophagidae (Diptera) of Fennoscandia and Denmark (*Faun. Ent. Scand.* **19**), 210; **Pitkin, B.R.,** 1988, Lesser Dung Flies Diptera Sphaeroceridae, (*Handbks Ident. Br. Insects*), 138; **Sheail, J.,** 1987, Seventy-Five years in Ecology: the British Ecological Society, 94; **Sims, R.W., Freeman, P. & Hawksworth, D.L.,** 1988, Key Works to the Fauna and Flora of the British Isles and North-Western Europe, 5th edn, 149; **Skou, P.,** 1986, The Geometroid Moths of North Europe, 94; **Smith, F.G.A.M.,** 1987, An Illustrated Catalogue of the Rothschild Collection of Fleas (Siphonaptera) in the British Museum (Natural History). VII, Malacopsylloidea (Malacopsyllidae and Rhopalopsyllidae), 176; **Stubbs, F.B.,** 1986, Provisional Keys to British Plant Galls, 122; **White, I.M.,** 1988, Tephritid Flies, Diptera Tephritidae, *Handbks Ident. Br. Insects*, 235.

Ringer for bathing termite (Isoptera) gut tissues, a simple, 233

Scotland: *Cyphon kongsbergensis* Munster (Col., Scirtidae) in Wester Ross, 156; *Gymnetron beccabungae* (L.) (Col., Curculionidae) in, 51; The status of *Noterus* spp. (Col., Noteridae) in, 71; Two Coleoptera (Lathridiidae, Derodontidae) records from Roxburghshire (VC80), 198

Scottish Highlands, two records of *Salidula opacula* (Zett.) (Hem., Saldidae) from the, 249

Secondary sexual character in the legs of male *Nesolynx glossinae* (Waterston) (Hym., Chalcidoidea, Eulophidae), a remarkable, 231

Skye, *Lamproplax picae* (Flor.) (Hem., Lygaeidae) on the Isle of, 182

Solomon Islands, notes on marine *Rheumatometroides* (Hem., Gerridae) from the, 63

South Africa, nesting behaviour and prey of two species of *Stizus* (Hym., Sphecidae, Nyssoninae), notes on, in, 157; observations on two species of Curculionidae (Col.) associated with cycad cones in, 129

Sudan, foraging of the harvester ant *Messor galla* Mayr. (Hym., Formicidae) in West, with some indication of pest potential, 207

Suffolk Sandings and Breckland, notable wasps and bees (Hym., Aculeata) in, 232

Surrey, *Brachygaster minutus* (Olivier) (Hym., Evaniidae) in, 84; *Miridius quadrivirgatus* (Costa) (Hem., Miridae) found inland in, 174

Synonymy, new, in rice-associated leafhoppers of the genera *Nephotettix* and *Cofana* (Hem., Homoptera, Auchenorhyncha, Cicadellidae), 135

"Travellers", Stratiomyid (Diptera), the larval habits of some, with new synonymy, 141

Troisième Conference Internationale des Entomologistes d'Expression Francaise, v

Wales, *Ceutorhynchus punctiger* Gyll. (Col., Curculionidae) in South, 30; *Euophryum confine* Broun (Col., Curculionidae) in South, 106

Warwickshire, *Agnocoris reclairei* (Wagner) (Hem., Miridae) in, 36

Wilderness Watch, xv

Windsor, *Corticaria linearis* (Payk.) (Col., Lathridiidae) at, 181

Worcestershire, an inland record of *Calathus mollis* (Marsham) (Col., Carabidae) in, 198; changing environments and Coleoptera faunas from Aston Mill, 187; *Harpalus dimidiatus* Rossi (Col., Carabidae) in, 118

Ynes Enlli (Bardsey I.), meniscus midges (Dipt., Dixidae) on, 118

Yorkshire, *Ectemnius sexcinctus* (F.) (Hym., Sphecidae) – a wasp new to, 250

SPECIAL INDEX

New genera, and species, etc., are indicated by heavy type, genera and species new to Britain by an asterisk (*).

ARACHNIDA

Acarus nidicolous, 237–9
Aranella cucurbitica, 50
Dicranopalpus ramosus, 236
harvestman, 48
Linyphia triangularis, 50
Mitopus morio, 50, 256
Philodromus aureolus, caespitum, dispar, 50
spider web, 92
Tegenaria, 91
Tetragnatha degeeri, 50
Theridion sisyphium, 36
Xysticus sp., 50

COLEOPTERA

? Coleoptera, 200
Acupalpus meridianus, 189, 197
Adalia decempunctata, 50
Agabus guttatus, 189
Agonum albipes, dorsale, fuliginosum, marginatum, moestum, muelleri, 188; obscurum, 118, 188
Alcides, 201
Aleochara bipustulata, curtula, 190
Allocorynidae, 131
Allocorynus, 129
Aloconota gregaria, 190
Alophus triguttatus, 192, 194
Altica lythri, 118, 191
Amalus scortillum, 128
Amara aenea, apricaria, aulica, bifrons, consularis, 188; convexiuscula, 79, 150; familiaris, 188; fulva, 188–197; montivaga, similata, tibialis, sp., 188
Amischa analis, 190
Amophocerus, 129–31
Anaspis garneysi, 191; maculata, rufilabris, 50
Anisodactylus binotatus, 79, 150
Anotylus inustus, rugosus, 189
Anthonomus, 100
Anthrenus sarnicus, 96, 98; verbasci, 98
Antliarrhinidae, 131
Antliarrinus zamiae, 129
Aphodius, 193; fossor, 118, 190; granarius, 190; lapponum, 4; obliteratus, 236; rufus, 190; sphacelatus, 190, 196; sp. 190
Apion curtirostre, 118; dichroum, 191; ervi, sicardi, 118
Araecerus, 131
Araucariini, 130
Arhopalus rusticus, 250
Aridius bifasciatus, 50
Arpedium brachypterum, 4
Artematopus, 210
Atheta fungi, 190
Atomaria, 187; pulchra, 252
Axinotarsus marginalis, 96, 99; pulicarius, 99
Badister bipustulatus, 189
Bembidion aeneum, 188, 194; articulatum, bipunctatum, dentellum, femoratum, genei, guttula, 188; iricolor, 150; obtusum, 188; punctulatum, 150; quadrimaculatum, tetracolum, 188
Bitoma crenata, 252
Blaps mucronata, 191
Bledius opacus, 189
Boganidae, 131
Bostrychid, 204
Bradycellus verbasci, 189
Bruchus rufipes, 252
Buprestid, 204
? buprestid, 203
Buprestidae, 131, 203–4
Byrrhus fasciatus, 191; sp., 191, 194
Calathus fuscipes, 188, 197; melanocephalus, 188, 194; mollis, 198
Calendra, 203
Calvia quattuordecimpunctata, 50
Cantharis lateralis, nigra, nigricans, 191
Carabus glabratus, nitens, 4; monilis, 187, 196; violaceus, 118, 187
Carpelimus corticinus, gracilis, 189; zealandicus, 150, 189, 197
Carpophilus ligneus, marginellus, 132
Cassida nebulosa, 168
Cateretidae, 131
cerambycid larva, 200
cerambycid larvae, 202
Cerambycidae, 202–4
Cercyon marinus, 150
Cerylon ferrugineum, histeroides, 252
Cetoniinae, 131
Ceutorhynchus euphorbiae, 117; marginatus, mixtus, pilosellus, punctiger, 30; querceti, 128
Chalcoides fulvicornis, 118
Chiloporata longitarsis, 190
Chlaenius vestitus, 189
Choleva angustata, 189, 197
Chrysobothris, 203
Cicindela campestris, 82
Cionus scrophulariae, 150
Cleonus piger, 192, 196
Clivina fossor, 118, 188
Coccinellidae, 191
Coelambus confluens, 150
Colymbetes sp., 189
Coccinella 5-punctata, 177–81; 11-punctata, 180
Colon dentipes, 242

Corticaria dubis, eppelsheimi, linearis, 181
Corticarina fuscula, 191; gibbosa, lambiana, similata, truncatella, 79
Cortinicara gibbosa, 50
Cossoninae, 130
Crepidodera ferruginea, transversa, 191
Criocephalus rusticus, 250
Cryptogenius, 210
Ctenicera pectinicornis, 191
curculionid larvae, 204
Curculionidae, 129–31, 203–4
Cyphon kongsbergensis, ochraceus, 156
Cytilus sericeus, 191, 194
Dactylispa discoidalis, 23
Demetrias atricapillus, 189
Diacheila arctica, 188, 194
Dicladispa, 23
Drusilla canaliculata, 190
Dryophilus pusillus, 252
Dyschirius aeneus, 150
Elaphrus cupreus, riparius, 188, 194; sp., 194
Enaretta castelnaudi, 200
Ephistemus globulus, 252
Euophryum confine, 106
Eusphalerum luteum, 252
Furcipus rectirostris, 96, 99–100
Galerucella lineola, 191
Geotrupes spiniger, stercorarius, sp., 190
Glischrochilus hortensis, 191
Glocianus, 30
Gnypeta carbonaria, 150
Grypus equiseti, 192, 194
Gymnetron antirrhini, 28; beccabungae, 25, 51; collinum, linearia, 28; lloydi, 25–6; marshalli, melas, 25, 27, 29; pascuorum, 25, 27, 28–9; plantaginis, 25, 28–9; veronicae, 25, 51
Harpalus affinis, 118, 188; ardosiacus, 188, 197; dimidiatus, 118; rubripes, 189; rufibarbis, 188, 197; rufipes, 118, 188
Helophorus sibiricus, 189–194; sp., 189
Henoticus serratus, 252
Heterocerus, 194; cf hispidulus 191, 194
Hispa armigera, 23; hispa, similis, 19
Hister quadrimaculatus, 189, 196; striola, 189
Homalota plana, 252
Hydrobius fuscipes, 189, 194
Hydrochara caraboides, 189, 195
Hydroporus planus, sp., 189
Hypnoidus rivularis, 191, 194
Ilybius sp., 189
Laccobius striatulus, 189
Laemostenus terricola, 188
Lamprohiza splendidula, 182
Lamprosoma concolor, 236
Lampyria noctiluca, 182
Languriidae, 131
Laricobius erichsonii, 198
Lathridius anthracinus, 198; consimilis, 79; minutus, pseudominutus, 198
Lathrobium elongatum, fulvipenne, sp., 190
Leiodes ferruginea, 189, 197
Leistus ferrugineus, fulvibarbis, 187; rufomarginatus, 18, 95–6; spinibarbis, 98
Leptura rufa, 182
Lissomus, 210
Longitarsus reichei, 118
Loricera pilicornis, 188
Lymexylonidae, 203
Malachius, 99; bipustulatus, 50
Mecinus pyraster, 192
Mecysolobus, 201
Megalopodinae, 131
Megascelinae, 131
Melanophthalma curticollis, 79; suturalis, transversalis, 140
Meloe rugosus, 127
Meloid larvae, 167
Megabletus obscuroguttatus, 189
Migdolus, 210
Monocheirus callicanthus, 19
Morychus aeneus, 191
Myrrha octodecimguttata, 50
Nebria brevicollis, 187
Nicrophorus humator, 189
Notaris acridulus, aethiops, bimaculatus, 192, 194; spp., 194
Noterus clavicornis, 71–2; crassicornis, 72
Notiophilus aquaticus, 187, 194; palustris, 187; substriatus, 188
Olibrus aeneus, affinis, corticalis, 191
Olophrum fuscum, 4, 189, 194
Omophron limbatum, 95–6
Oncomera femorata, 236
Onthophagus joannae, 190, 196
Ostoma ferrugineum, 96, 98–9
Otiorrhynchus, 197, arcticus, clavipes, ligneus, nodosus, rugifrons, 192, 194; scaber, 132; singularis, 50; sulcatus, 192
Oxytelus lacqueatus, 50
Ozoneus prolixus, 202
Parabathyscia wollastoni, 50
Paralister purpurascens, 189, 197
Pediacus dermestoides, 252
Phaedon cochleariae, 191
Pharaxonothinae, 131
Phausis splendidula, 182
Philonthus intermedius, 118; politus, 190; rotundicollis, 244; varians, 190
Phloeophagus hipidus, 129
Phoracantha semipunctata, 201
Phrixothrix, 210
Phyllobius roboretanus, 192
Phyllodecta cavifrons, laticollis, vitellinae, vulgatissima, 24
Phyllopertha horticola, 190, 197
Phytobius quadrituberculatus, 192
Phytodecta decemnotata, olivacea, 17; pallida, viminalis, 17–8
Plateumaris sericea, 191, 195
Platystethus nodifrons, 189
Porculus, 210
Porthetes zamiae, 129–31
Pseudotriphyllus suturalis, 252

Psylliodes picina, 191
Pterostichus adstrictus, 188, 194; aethiops, 118; cupreus, 118, 188; diligens, 118; macer, 118; madidus, melanarius, niger, 188; nigrita, 118; strenuus, 118, 188; vernalis, 188; sp., spp., 194
Ptilinus pectinicornis, 252
Ptiliolum caledonicum, 138
Ptilium, 187; affine, exaratum, horioni, incognitum, 138
Quedius aridulus, cinctus, 190; mesomelinus, 236; molochinus, picipes, schatzmayri, tristis, sp., 190
Rhagonycha fulva, 191
Rhinus (s.g.), 28
Rhopalotria, 129
Rhynchaenus pascuorum, 17; sparsa, 29; testaceus, 167
Rugilus orbiculatus, 190
Sciaphilus asperatus, 192
Sciodrepoides watsoni, 189
Scolytidae, 203
Scotocryptodes, 210
Sepedophilus immaculatus, 12; marshami, 12, 190; testaceus, 12
Sermylassa halensis, 236
Sibinia potentillae, 25–6; primitus, 25, 28
Silpha obscura, 189, 196
Simplocaria semistriata, 190, 194
Sitona lineatus, 192
Sitophilus, 203
Sphenophorus phaenicensis, 203
Sphenoptera, 203; gossypii, 199, 203; laticollis, 203
Staphylinus ater, globulifer, 190; olens, 79, 190
Stenus biguttatus, canaliculatus, clavicornis, 189; crassus, formicetorum, 137; juno, 189; melanopus, 137, 150; subaeneus, 50
Stephostethus lardarius, 50
Sunius propinquus, 190
Synuchus nivalis, 188
Tachinus jacuticus, 190, 194
Tachyporus hypnorum, obtusus, 190; pallidus, 118
Tachyusa constricta, 190
Tesserocerus, 210
Thanatophilus dispar, 189, 194; latericarinatus, 194; rugosus, 189, 197
Tomoxia biguttata, bucephala, 252
Trachyphloeus bifoveolatus, scaber, 132
Trechus micros, obtusus, quadristriatus, 188; rivularis, 4
Tryphocharia sp., 201
Tychus niger, 190
Xantholinus jarrigei, linearis, longiventris, 190
Xyleborus saxeseni, 252
Xystrocera festiva, 203

COLLEMBOLA

Entomobrya nivalis, 49

DERMAPTERA

Carcinophora americana, 13–4, 16; **discalis**, festiva, haenschi, 14, 16; **lateralis**, percheroni, rosenbergi, 13–4, 16
Euboriella annulata, janeirensis, stali, 16
Pyragropsis thoracica, 16

DIPTERA

Acinoeuphranta, 31
Acritochaeta (s.g.), 73, 142
Acrostilpna similata, 219
Aedes aegypti, 122; domesticus, furcifer, 115
Agromyza buhriella, 83; pagana, paganella, 84; rufipes, spiraeoidearum, s. arunci, 83
Agromyzidae, 7, 62
Amiota leucostoma, 52
Anapausis **aberdarica**, 39; absoluta, 37–40; conspicua, 41; **elgonensis**, 39; irritata, 41; **kashmirica**, 40–2; **stapediformis**, 41–3; truncata, 41; **zealandica**, 41–2
Anopheles maculipennis, 52
Anthomyia *confusanea, lituratа, 211–3; pullula, 211
Anthomyzidae, 7
Aphanotrigonum trilineatum, 243
Apocephalus laceyi, 171, 173; wallerae, 171
Argyra elongata, 243
Aricia moriens, 218–9
Asphondylia matabai, 201
Atherigona addita, 76; aurifacies, 78; **chirinda**, 75–6; **decempilosa**, 76–8; haplopyga, 75; **maliensis**, 73–5; mambillaensis, perfida, 75
Auxanommatidia myrmecophila, 173
Bibio sp., 50
Bibionidae, 170
Bicellaria subpilosa, 243
Botanophila, 219; apiciseta, flavisquama, 217–8; **fonsecai**, 213–7; gnavoides, gnavula, 218; *helviana, 217; jacobaeae, 217–8; moriens, 218; pseudomaculipes, 217; setosa, 216; spinosa, 218; trigemina, 213, 215–6; trigeminata, 215–6; trigeminoidea, 216; trigeminoides, 215–7
Calliphoridae, 7
Camillidae, Camilla sp., 7
Campsicnemus compeditus, 243
Carnidae, Carnus hermapterus, 7
'Cecidomyiid', 201
Chamaemyiidae, 62

Cheilosia illustrata, 101
Chirosia similata, 219
Chloropidae, 7
Chrysochlora haterius, plana, vespertilio, 142
Chrysops caecutiens, 122
Chyromyidae, 7
Cochliomyia hominivorax, 62
Conopidae, 170
Craspedochoeta, 212–3; luteipennis, 219; pullula, 211
Culex modestus, molestus, 52; pipiens, 52, 107, 115; torrentium, urichii, 115
Culiseta longiareolata, 52
Cyclopsia, 31
Cylindrotoma distinctissima, 8
Dasyphora cyanella, 7
Dilophus febrilis, 50
Dimecoenia tristanensis, 62
Diplonevra mortimeri, watsoni, 173
Dixella aestivalis, 107; amphibia, 107, 112–6; attica, 118; autumnalis, 107, 112–3, 116; californica, 107, 114; filicornis, 107, 112–3, 116; martinii, 118
Drosophila acuminata, ananassae, iri, polychaeta, 52
Drosophilidae, 141; indet, 45
Dufouria, 62
Egle bicaudata, inermis, 220; *subarctica, 219–20
Empis verralli, 243
Episyrphus balteatus, 132
Eriozona syrphoides, 97, 100–1, 132
Eristalis horticola, 36
Ernoneura argus, 230
Euphranta, 31
Eustalomyia hilaris, 97, 101
'fly gall', 201
gout fly, 62
Heleomyzidae, 7
Hermetia illucens, 141–2
Hylemyia, 220; apiciseta, 217; nudithorax, spinosa, 218
Ichneumonosoma, 31
Indophranta humerata, 31–3
Iridophora clarki, 171
Leopoldius brevirostris, 153–6; signata, signatus, 153–5
Leucophenga paracapillata, 45
Leucophora, 216
Liriomyza pisivora, 83
Lycoriella agraria, 7
Macrocera fasciata, 91
Medetera impigra, pinicola, 174
Medina collaris, 139; luctuosa, 139–40; melania, multispina, 139; *separata, 139–40
Megaselia benebarbata, 48; caudifera, 46; chaetogaster, 48; divergens, 186; **falklandensis**, 183–6; kodongi, 170–1; **malvinensis**, 184–6; masatierrana, 185–6; nigricauda, 46; persecutrix, 169–73; plaumanni, 186; rufipes, 48; sembeli, 169–171; similis, 185; **termitomyia**, 46–7; teutoniae, 186; zariaensis, 46–8
Melina luteipennis, 219
Meliniella bisinuata, griseifrons, sikisima, watanabii, 219
Microselia texana, 171
miltogrammine flies, 166–7
Minella, 62
Molophilus ater, 243
Muscidae, 7
Myrmosicarius texanus, 171
Nematocera, 170
Neoascia podagrica, 50
Neoexaireta spinigera, 141
Neottiophilidae, Neottiophilum praeustum, 7
Norellia spinipes, tipularia, 105
Ophiomyza alliariae, 83
Ophyra chalcogaster, 45
Opifex fuscus, 115
Paregle *atrisquama, 220; audacula, 221
Pegohylemyia, 217–20; flavisquama, 218
Pegomya, 219; betae, 224; circumpolaris, 224–6, 228–9; cunicularia, 224; *depressiventris, 222–4; dulcamarae, 223; exilis, 224; fulgens, 223–8; furva, 224–8; geniculata, 224; genupuncta, 223; hyoscyami, laticornis, steini, 223–4; tabida supersp, 224; tenera var. obscurior, 228; *transgressa, 223–8
Phaonia consobrina, 243
Phlebotomus papatasi, 122
Phora greenwoodi, 81–2; speighti, 81
Phytomyza anthyllidis, biseta, 83; kasi, latifrons, 84; mylini, podagrariae, 83; pulchella, pulchelloides, 84
Piophilidae, 141
Plastophora, 45; divergens, 186
Platypalpus alter, 243; longimana, 253
Pliomelaena **quadrimaculata**, 33–5
Protocalliphora azurea, 7
Pseudacteon brevicauda, 171, 173; californiensis, 171
Pseudomyopina, 219; moriens, 218
Psorophora confinnis, 115
Pterogramma, 45
Ptilona, 31
Rhamphomyia marginata, 97, 100
Sarcophaga carnaria, 210
Sargus, 142
Scaptomyza griseola, montana, trochanterata, 52
Scatopsidae, 7
Scenopinidae, 141
Sciaridae, 7
Silba aristella, 62
Simulium, 122; rubrithorax, 44
Speolepta leptogaster, 89–93
Spelobia talparum, 7
Sphaeoceridae, 7, 138
Spilogona litorea, longipes, 244; *trigonata, 243–4

Stomoxys calcitrans, 122
Stylogasterine, 170
Syrphidae, 7, 62
Tachinidae, 7, 62
Tephritidae, 7, 235
Terellia **uncinata**, 53–7, 59–60; vectensis, 54; virens, 53–60; zerovae, 53–5, 59–60
Thaumatomyia notata, 7
Thricops albibasalis, cunctans, hirsutulus, 243
Tipula oleracea, paludosa, 123–6; subnodicornis, 243
Tripeta syllibi, 56
Trypeta virens, 56
Xanthocnemia, 217

HEMIPTERA

Agnocoris reclairei, 36
Aleyrodes lonicerae, proletella, 72
Atractotomus magnicornis, 49
Cinara fresai, 49; pinea, 101
Cofana unimaculata, 136–7
Deltocephalus trimaculatus, 136–7
Dichrooscytus valesianus, 49
Hynesionella, 71
Lamproplax picea, 182
Miridius quadrivirgatus, 174
Myrmedobia tenella, 49
Naboandelus, 71
Neanoecia krizusi, 4
Nephotettix malayanus, nigropictus, 135–6; olivacea, 136; oryzii, 135–6; parvus, 136; virescens, 135–6
Orsillus depressus, 241–2; maculatus, reyi, 241
Phrynomorphus olivacescens, 135, 137
Placochilus seladonicus, 205
Rhopalosiphum padi, 17
Rheumatometroides **aqaaqa**, 64–5, 69–70; browni, 63–70; **matawa**, 64, 67–70
Saldula opacula, saltatoria, 249
Selenocephalus virescens, 135
Stenobates, 63, 69, 71; biroi, 70; insularis, 69, 71
Temnostethus gracilis, pusillus, tibialis, 206
Tettigonia unimaculata, 136
Trepobatinae, 71

HYMENOPTERA

Abia sericea, 127
Acantolyda posticalis, 127
Andrena hattorfiana, 168
Anophonyx destructor, 62
Antiolcus apicalis, 120
Argogorytes fargeii, 232
Aspidibracon hespirivorus, noyesi, pierri, 200
Atanycolus, 120
Atanycolus-group, 202, 204
Bathyaulax, 199; spp., 200–201
Batozonellus fuliginosus, 158
Bembecinus, 166
Brachygaster minutus, 84
Bracon, 10, 199; agrili, 120; crassitarsis, 119; hispae, 23; kirkpatricki, 201; sanguineus, 119; thalessiformis, 120; spp., 201
Braconina, 202
Braconella, 199; sp., 201
Callibracon limbatus, 201
Callirhytis erythrocephala, 50
Camponotus gigas, 169; rufipes, 173
Campyloneurus asphondylai, 201; busckii, 120; ceylonicus, sp., 201
Cephalcia lariciphala, 62, 127
Cervellus, 199; sp., 202
Coeloides, 9
Colletes marginatus, 232
Compsobracon, 10, 119, 199; sp., 202
Crabro cribrarius, 250; scutellatus, 232
Cratocnema spp., 202
Cyclaulax, 199
Cyanopterus diversus, peculiaris, 120; sanguineus, 119
Digonogastra, 119, 199; agrili, augustus, busckii, carapunae, diversus, fortis, reduvioides, starski, wolcottii, 120; spp., 202
Dolerus bimaculatus, ferrugatus, 127; gessneri, 168
Dolichovespula media, 97, 102, 106; m. geeri, 106; norwegica, 104–5, 117, 133–4, 143–6, 148; *saxonica, 97, 101–2, 103–5; sylvestris, 103–5, 117, 133–4, 143–8
Ectemnius continuus, 250; rubicola, 232; sexcinctus, 250
Empria pumila, 127
Encarsia inaron, tricolor, 72
Esenga, 20–1
Euderomphale *chelidonii, 72
Euurobracon cephalotes, 203
Euvipio, 199, 200
Formica rufa, 137
Glyptomorpha andrieui, desectus [desertus], desertor, spp., 203
Gorytes tumidus, 232
Habrobracon, 199
Halictus confusus, 232
Hemibracon abunensis, apicalis, 120
Heterospilus koebelei, 202
Iphiaulax, 202; abunensis, agrili, augustus, carapunae, fortis, peruensis, reduvioides, 120; sanguineus, 119; starski, wolcottii, 120
Iridomyrmex, 209; cordatus, 169
Isomecus spp., 203
Lasioglossum brevicorne, 232

Lasiophorus polaris, semirufus, 121
Lasius alienus, flavus, fuliginosus, meridionalis, niger, 4
Lindenius panzeri, 2323
Loderus eversmanni, pratorum, p. gilvipes, vestigialis, 168
Macroneuroides erythropleura, 120
Macrophya montana, 127
Megabracon erythropleura, peculiaris, thalessiformis, 120
Megacoeloides annulicornis, 9–12
Messor galla, 207–9
Microbracon hispae, 20;
Microdiprion pallipes, 127
Monogastra agrili, augustus, wolcottii, 120
mutillids, 167
Myosoma, 199; chilonus, chinensis, spp., 204
Myrmica sabuleti, 4
Nematus umbratus, 127
Nesolynx albiclavus, 231; glossinae, 231–2; seyrigi, 232
Odontoscapus, 199; spp., 204
Odynerus simillimus, 206
Ophthalmobracon (s.g.), 201
Osmia inermis, 102; uncinata, 97, 102
Oxybelus argentatus, mandibularis, 232
Pachynematus imperfectus, moerens, 127
Palabracon polaris, semirufus, 121
Pamphilius balteatus, 127
Paravespula, 103; germanica, 143–5, 148; vulgaris, 143–6, 148
Periclista pubescens, 127
Philanthus triangulum, 232
Physaraia sp., 204
Plesiobracon, 20
Podalonia affinis, 232
Pristiphora compressa, laricis, 62
Pseudoshirakia yokahamensis, 204
Pseudovipio sp., 204
sawfly larvae, 203
Scutibracon hispae, 19–23
Sharliphora amphibola, 62
Siricidae, 203
Stenobracon, 199, 200
Stizus atrox, 164; chrysorrhoeus, 163; dewitzi, 164; distinguendus, 157, 166; fuscipennis, 157–9, 161–6; imperialis, 157, 160–7; iridis, marshalli, 157; pentheres, 164; pulcherrimus, ruficornis, 157
Syntormophyrum glossinae, 231
Triaspidogastra, 199; sp., 205
Vespa crabro, 102, 106, 143–5
Vespula rufa, 143–8; vulgaris, 134
Vipio, 203; sanguineus, 119

ISOPTERA

Macrotermes michaelseni, 233–4
Odontotermes magdaleneae, 45
Zootermopsis nevadensis, 233–4

LEPIDOPTERA

Acidion socium, 202
Amatidae, 45
Apisa bourgognei, 45
Chilotraea auricilia, 201
Coleophora, 201
Cupido argiades, 80
Diatraea, 121; saccharalis, 203
Doratoperas atroparsellus, 202
Drepanidae, Drepaninae, 94
Erionota thrax, 200
Eupithecia sobrinata, 49; spp., 94
Gelechoidea, 201
Geometridae, Geometroidea, 94
Gnorimoschema heliopa, 201
Hypsipyla grandella, 203
Idaea, 94
larva, 204
lepidopteran, 236
lepidopteran pupa, 200
Manduca secta, 239
Melitaea phoebe, 80
Parategetica, 202
Parnara mathias, 200
Paspalum fasciculatum, 204
Pectinophora gossypiella, 201
Pelopidas mathias, 200
Platyedra, 201
Recurvaria, 201
Scirpophaga nivella, 204
Syntomidae, 45
Tegeticula yuccasella, 202
Thytirinae, 94
Topeutis sp., 204
Yponomeuta evonymellus, 17

MANTODEA

mantid nymphs, 158
Phyllocrania paradoxa, 159
Tarachodes, 159

MECOPTERA

Panorpa communis, 36

NEUROPTERA

Chrysopa cf flavifrons, 49
Conwentzia psociformis, 49
Hemerobius lutescens, 49; marginatus, 82

ODONATA

Somatochlora metallica, 151–2; uchidae, 152

ORTHOPTERA

Acrididae, 161
Cycloptiloides, 45
Eremidium, sp., 161
Gastrimargus, 161
Hoplolopha sp., 161
Lentula sp., Lentulidae, 161
Oedaleus, 161
Pamphagidae, 161
Pyrgomorpha sp., Pyrgomorphidae, 161

PHTHIRAPTERA

Lutridia, 248; exilis, 245–8; lutrae, matschiei, 247
Neolutridia, 247–248
Polyplax serrata, 246
Trichodectes canis, 245

PSOCOPTERA

Caecilius sp., 49
Cerobasis guestfalica, 49

SIPHONAPTERA

Archaeopsylla erinacei erinacei, 237–9

THYSANURA

Dilta hibernica, 48

GROUPS OTHER THAN INSECTA AND ARACHNIDA

Amphibia, 196
Aves, 5–7, 196
Chilopoda, 48, 91
Isopoda, 48, 79, 91
Mammalia, 5–7, 195–7, 237–9, 245–8
Mollusca, 48, 79, 91, 193, 195–6, 236

ERRATA AND CORRIGENDA

Page	Line	For	Read
49	3	(Hemiptera: Miridae) Meyer-Dür	Meyer-Dür (Hemiptera: Miridae)
50	3 (up)	*arureolus*	*aureolus*
62	15	Lonchaeidne	Lonchaeidae
62	18 (up)	*lariciphila*	*lariciphala*
"	13 (up)	"	"
72	13 (up)	*chelidinii*	*chelidonii*
83	2 (up)	Görschke	Gröschke
203	22	*desectus*	*desertus* (Bradley)
v	14	Josifon	Meyer-Dür

VOL **124**

102	20 (up)	TQ 715343	TQ 636384

Printed in England by V.B. Pike Printers, Kettering, Northamptonshire